



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Classification of plants.**—A third edition of Professor BESSEY's *Outlines of plant phyla* has appeared. The previous editions were noted in this journal (51:317. 1911; 53:275. 1912) and the general features of the classification given. In the new edition the 14 phyla remain the same, but the families have increased from 648 to 652. The relation of the conventional four great groups to this scheme may be indicated by the statement that the thallophytes are broken up into 7 phyla, the bryophytes remain as a single phylum, the pteridophytes become 3 phyla, and the spermatophytes 3.—J. M. C.

**Perithecium of Polystigma.**—BLACKMAN and WELSFORD<sup>37</sup> have investigated the development of the perithecium of *Polystigma*, and have discovered that while well marked ascogonia occur, they disorganize without producing ascogenous hyphae, and that the spermatia are also functionless. It differs from most of the Ascomycetes in which the normal sexual process is absent by the fact that both sex organs are distinctly produced, but that both are abortive.—J. M. C.

**Wound reactions in fern petioles.**—HOLDEN<sup>38</sup> induced wound reactions in the petioles of 37 species of ferns, the wounds being thin superficial shavings made with a scalpel in three regions: the curled apical portion, the region of pinna insertion, and the region below pinna insertion. Various reactions were obtained, which differ too much in details to be summarized in a review, but all of which are interesting contributions to the subject of wound reactions.—J. M. C.

**Mycorrhiza of Asarum.**—SCHWARTZ<sup>39</sup> has studied the mycorrhiza of *Asarum europaeum*, finding it limited to the cortical region abutting on the steles of young roots. Thick-walled swellings were found on some of the hyphae, representing a resting stage.—J. M. C.

**Lateral archegonia in Pinus.**—SAXTON<sup>40</sup> has made the very interesting discovery of a female gametophyte of *Pinus maritima* which bears two lateral groups of two archegonia each, and no terminal (micropylar) archegonia at all.—J. M. C.

<sup>37</sup> BLACKMAN, V. H., and WELSFORD, E. J., The development of the perithecium of *Polystigma rubrum* D.C. Ann. Botany 26:761-767. pls. 70, 71. 1912.

<sup>38</sup> HOLDEN, H. S., Some wound reactions in filicinean petioles. Ann. Botany 26:777-793. pls. 73, 74. fig. 1. 1912.

<sup>39</sup> SCHWARTZ, E. J., Observations on *Asarum europaeum* and its mycorrhiza. Ann. Botany 26:769-776. pl. 72. 1912.

<sup>40</sup> SAXTON, W. T., Note on an abnormal prothallus of *Pinus maritima* L. Ann. Botany 26:943-945. fig. 1. 1912.